




CD152 [CTLA-4] non-lytic Recombinant Protein

Cat. No.: 90-413




Ψ Specifications

SPECIES:	Human
SOURCE SPECIES:	CHO cells
SEQUENCE:	The extracellular domain of human CD152 [CTLA-4] (aa 37-160) is fused to the N-terminus of the Fc region of a mutant human IgG1.
FUSION TAG:	Fc Tag
TESTED APPLICATIONS:	
APPLICATIONS:	This recombinant proteins is for research use only.

Ψ Properties

PURITY:	≥98% (SDS-PAGE). Endotoxin level is less than 0.06EU/ µg protein (LAL test; Lonza).
PHYSICAL STATE:	Lyophilized
BUFFER:	Lyophilized from 0.2µm-filtered solution in PBS. Reconstitute at 100 µg/ml in sterile PBS.
STORAGE CONDITIONS:	Stable for at least 1 year after receipt when stored at -20 °C. Working aliquots are stable for up to 3 months when stored at -20 °C.

OFFICIAL SYMBOL:	CTLA4
ALTERNATE NAMES:	CTLA-4, Cytotoxic T-lymphocyte Protein 4
ACCESSION NO.:	NP_005205
PROTEIN GI NO.:	21361212
GENE ID:	1493

 Background and References

BACKGROUND:	<p>CD152 (CTLA-4) and CD28, together with their ligands B7-1 and B7-2, constitute one of the dominant costimulatory pathways that regulate T and B cell responses. CD152 and CD28 are structurally homologous molecules that are members of the immunoglobulin (Ig) gene superfamily. Both CD152 and CD28 are composed of a single Ig V-like extracellular domain, a transmembrane domain and an intracellular domain. CD152 and CD28 are both expressed on the cell surface as disulfide-linked homodimers or as monomers. CD152 was originally identified as a gene that was specifically expressed by cytotoxic T lymphocytes. However, CD152 transcripts have since been found in both Th1 and Th2, and CD4+ and CD8+ T cell clones. Whereas, CD28 expression is constitutive on the surfaces of 95% of CD4+ T cells and 50% of CD8+ T cells and is down regulated upon T cell activation, CD152 expression is upregulated rapidly following T cell activation and peaks approximately 24 hours following activation. Although both CD152 and CD28 can bind to the same ligands, CD152 binds to B71 and B72 with 20-100-fold higher affinity than CD28.</p>
--------------------	---

ANTIBODIES FOR RESEARCH USE ONLY.

For additional information, visit ProSci's [Terms & Conditions Page](#).