




SARS-COV-2 Spike RBD Protein Antibody [2F7A11]

Cat. No.: 24-040



Ψ Specifications

HOST SPECIES:	Mouse/Human chimeric
SPECIES REACTIVITY:	Virus
IMMUNOGEN:	Hek293 derived SARS-COV-2 SPIKE RBD Arg319-Phe541 Accession #YP_009724390.1
TESTED APPLICATIONS:	ELISA
APPLICATIONS:	SARS-COV-2 Spike RBD Sandwich Elisa: ELISA Capture: Recommended Concentration:1-4ug/mL, Sample: SARS-CoV-2 Spike RBD Protein Antibody (CAP) ELISA Capture: Recommended Concentration:0.05-0.2ug/mL, Sample: SARS-CoV-2 Spike RBD Protein Antibody Standard:Recommended Concentration: 15.6-1000pg/mL, Sample: Recombinant SARS-COV-2 Spike RBD Protein with His tag
SPECIFICITY:	SARS-COV-2

Ψ Properties

PURIFICATION:	Affinity purification < 1.0 EU/ug of the protein by LAL method.
CLONALITY:	Polyclonal

ISOTYPE:	IgG
CONJUGATE:	Unconjugated
PHYSICAL STATE:	Lyophilized
BUFFER:	Lyophilized from a 0.2 um filtered solution in PBS, pH7.4 with antibody at a final concentration of 1 mg/ml.
CONCENTRATION:	1 mg/ml
STORAGE CONDITIONS:	2 °C - 8 °C for one month, -20 °C to -80 °C for one year. Avoid repeated freeze-thaw cycles

Additional Info

OFFICIAL SYMBOL:	S
ALTERNATE NAMES:	SARS-CoV-2 (COVID-19, 2019-nCoV) Spike Antibody: Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), Surface Glycoprotein, Spike protein
ACCESSION NO.:	YP_009724390.1
GENE ID:	43740568
USER NOTE:	Optimal dilutions for each application to be determined by the researcher.

Background and References

BACKGROUND:	<p>The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that SARS-CoV-2 (COVID-19 coronavirus, 2019-nCoV) can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.</p>
--------------------	---

ANTIBODIES FOR RESEARCH USE ONLY.

For additional information, visit ProSci's [Terms & Conditions Page](#).