



SARS-CoV-2 (COVID-19) Spike RBD Monoclonal Antibody (B001N)

Cat. No.: 10-552




Figure 1 Western Blot Validation with Recombinant Protein

Loading: 10µg of recombinant protein per lane. Lane 1: 10-008 and Lane 2: 10-011. Antibodies: SARS-CoV-2 (COVID-19) Spike-ECD/RBD Monoclonal, 10-552, 1:1000. Secondary: Goat ant

ELISA experiment

	1	2	
A	2.347	2.399	:1:1000
B	2.036	2.216	:1:10000
C	2.287	2.139	:1:20000
D	2.441	2.624	:1:40000
E	2.104	2.292	:1:80000
F	1.784	1.713	:1:160000
G	1.186	1.213	:1:320000
H	0.105	0.086	Blank control

Figure 2 ELISA Test

Coating original concentration: 2 µg/mL, 100 µL/well samples are column 1: SARS-CoV-2 (COVID-19) Spike-ECD Recombinant Protein, 10-011, and column 2: SARS-CoV-2 (COVID-19) Spike-RBD Recombinant Protein, 10-008.
Antibodies: SARS-CoV-2 (COVID-19) Spike-ECD/RBD Monoclonal Antibody, 10-552.
Secondary: Goat anti-human IgG HRP conjugate at 1:10000 dilution.
Develop: 15min, 100 µL/well.
Stop: Stop buffer 50 µL/well.


Ψ Specifications

HOST SPECIES:	Human
SPECIES REACTIVITY:	Virus
IMMUNOGEN:	S-RBD recombinant protein
TESTED APPLICATIONS:	ELISA, Neut, WB
APPLICATIONS:	ELISA: 1:320000; WB: 1:1000~2000

PURIFICATION:	Greater than 95% as determined by reducing SDS-PAGE.
CLONALITY:	Monoclonal
CONJUGATE:	Unconjugated
PHYSICAL STATE:	Liquid
BUFFER:	PBS, pH7.2
CONCENTRATION:	2 mg/mL
STORAGE CONDITIONS:	Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.



OFFICIAL SYMBOL:	S
ALTERNATE NAMES:	SARS-CoV-2 (COVID-19, 2019-nCoV) Spike Antibody; Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), Surface Glycoprotein, Spike protein
ACCESSION NO.:	QHD43416.1
GENE ID:	43740568
USER NOTE:	Optimal dilutions for each application to be determined by the researcher.



BACKGROUND:	SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as 2019-nCoV (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. SARS CoV-2 Spike Protein derived from 2019-nCoV is composed of S1 domain and S2 domain. The S1 contains a receptor-binding domain (RBD) that can specifically bind to angiotensin-converting enzyme 2 (ACE2), the receptor on target cells.
--------------------	---

ANTIBODIES FOR RESEARCH USE ONLY.

For additional information, visit ProSci's [Terms & Conditions Page](#).