



# TXNDC4 Peptide

Cat. No.: 46-542P



## Ψ Specifications

<b>SPECIES:</b>	Human
<b>TESTED APPLICATIONS:</b>	Blocking
<b>APPLICATIONS:</b>	TXNDC4 Peptide is used for blocking the activity of TXNDC4 antibody (catalog number 46-542). The TXNDC4 antibody ships at 0.5 mg/mL and the TXNDC4 Peptide ships as a 100 ug pellet. When peptide is reconstituted in 200 uL water, the concentration will be 0.5 mg/mL, identical to that of the antibody. The best starting ratio would be 1:1 (which means molar excess of peptide relative to antibody when identical volumes are mixed). Mix equal volume of peptide and antibody at the required dilution and leave at ambient temperature for 20 minutes. Then incubate and develop in parallel two blots: one with the antibody and one with the antibody which has been pre-absorbed with the provided peptide.

## Ψ Properties

<b>PHYSICAL STATE:</b>	Lyophilized
<b>BUFFER:</b>	TXNDC4 peptide is supplied as 100 µg of dried peptide. When the peptide is reconstituted in 200 µL water, the concentration will be 0.5 mg/mL.
<b>STORAGE CONDITIONS:</b>	Store at -20 °C. When reconstituted minimize freezing and thawing.

<b>OFFICIAL SYMBOL:</b>	ERP44
<b>ALTERNATE NAMES:</b>	PDIA10, TXNDC4, KIAA0573, UNQ532/PRO1075, Endoplasmic reticulum resident protein 44, Thioredoxin domain-containing protein 4, ER protein 44
<b>ACCESSION NO.:</b>	NP_055866.1
<b>PROTEIN GI NO.:</b>	52487191
<b>GENE ID:</b>	23071

---

**ANTIBODIES FOR RESEARCH USE ONLY.**

For additional information, visit ProSci's [Terms & Conditions Page](#).